Flat Space Amplitudes and Conformal Symmetry of the Celestial Sphere

Sabrina Pasterski, ¹ Shu-Heng Shao, ² and Andrew Strominger ¹

¹Center for the Fundamental Laws of Nature, Harvard University, Cambridge, MA 02138, USA ²School of Natural Sciences, Institute for Advanced Study, Princeton, NJ 08540, USA

Abstract

The four-dimensional (4D) Lorentz group $SL(2,\mathbb{C})$ acts as the two-dimensional (2D) global conformal group on the celestial sphere at infinity where asymptotic 4D scattering states are specified. Consequent similarities of 4D flat space amplitudes and 2D correlators on the conformal sphere are obscured by the fact that the former are usually expressed in terms of asymptotic wavefunctions which transform simply under spacetime translations rather than the Lorentz $SL(2,\mathbb{C})$. In this paper we construct on-shell massive scalar wavefunctions in 4D Minkowski space that transform as $SL(2,\mathbb{C})$ conformal primaries. Scattering amplitudes of these wavefunctions are $SL(2,\mathbb{C})$ covariant by construction. For certain mass relations, we show explicitly that their three-point amplitude reduces to the known unique form of a 2D CFT primary three-point function and compute the coefficient. The computation proceeds naturally via Witten-like diagrams on a hyperbolic slicing of Minkowski space and has a holographic flavor.

Contents

1	Introduction	1
2	Conformal Primary Wavefunctions	3
	2.1 Integral Representation	4
	2.2 Analytic Continuation and the Massless Wavefunction	6
3	Massive Three-Point Amplitude	7
Α	Klein-Gordon Inner Product	10

1 Introduction

Quantum field theory (QFT) scattering amplitudes in four-dimensional (4D) Minkowski space are typically expressed in terms of asymptotic plane wave solutions to the free wave equation. Translation invariance is manifest because the plane waves simply acquire phases which cancel due to momentum conservation. The $SL(2,\mathbb{C})$ Lorentz invariance is more subtle as plane waves transform non-trivially into one another. In this paper we find a basis of $SL(2,\mathbb{C})$ primary solutions to the massive scalar wave equation and recast certain 4D scattering amplitudes in a manifestly $SL(2,\mathbb{C})$ covariant form. This form is very familiar from the the study of two-dimensional (2D) conformal field theory (CFT), in which $SL(2,\mathbb{C})$ acts as the global conformal group. The appearance of the 2D conformal group is no coincidence, since the Lorentz group acts as the global conformal group on the celestial sphere, denoted by \mathcal{CS}^2 , at null infinity where the asymptotic states are specified.

Studies of the $SL(2,\mathbb{C})$ properties of scattering amplitudes date back to Dirac [1], but a new reason has arisen for interest in the topic. When gravity is coupled, it was conjectured in [2–5] that the $SL(2,\mathbb{C})$ is enhanced to the full infinite-dimensional local conformal (or Virasoro) group. This conjecture was recently proven [6–8] to follow at tree level¹ from the new subleading soft graviton theorem of [15].² While the full Virasoro appears only when gravity is coupled, the scattering amplitudes of any QFT that can be coupled to gravity are

¹The subleading soft theorem has a one-loop exact anomaly [9–12] whose effects remain to be understood but are recently discussed in [13,14].

²One may hope that ultimately 4D quantum gravity scattering amplitudes are found to have a dual holographic representation as some exotic 2D CFT on \mathcal{CS}^2 , but at present there are no proposals for such a construction.

constrained to be 'Virasoro-ready'.³ This suggests that they should resemble a subset of 2D CFT correlators, likely involving complex and continuous conformal dimensions. Indeed it has already been observed [16,17] that soft photon amplitudes take the form of a 2D current algebra. Here we seek to understand the 2D description of 4D scattering amplitudes away from the soft limit.

This paper considers massive scalar three-point functions, and recasts them in the standard form of 2D CFT correlators on the celestial sphere \mathcal{CS}^2 . To summarize the result, let X^{μ} ($\mu = 0, 1, 2, 3$) be the flat coordinates on Minkowski space. A natural coordinate on \mathcal{CS}^2 where $X^{\mu}X_{\mu} = 0$ is

$$w = \frac{X^1 + iX^2}{X^0 + X^3} \,. \tag{1.1}$$

Lorentz transformations act as the global conformal group on \mathcal{CS}^2

$$w \to \frac{aw+b}{cw+d} \,. \tag{1.2}$$

Here the complex parameters a, b, c, d obey ad - bc = 1. We will construct a three-parameter family of solutions labelled by a point w on \mathcal{CS}^2 and an $SL(2,\mathbb{C})$ weight Δ (rather than the three components of spatial momenta \vec{p} which label plane waves) which transform as conformal primaries. We will find below they are naturally displayed in a hyperbolic slicing of Minkowski space, in harmony with the form of flat space holography advocated in [2]. $SL(2,\mathbb{C})$ then implies that the 4D tree amplitude takes the form

$$\tilde{\mathcal{A}}(w_i, \bar{w}_i) = \frac{C}{|w_1 - w_2|^{\Delta_1 + \Delta_2 - \Delta_3} |w_2 - w_3|^{\Delta_2 + \Delta_3 - \Delta_1} |w_3 - w_1|^{\Delta_3 + \Delta_1 - \Delta_2}},$$
(1.3)

where the 'OPE coefficients' C depends on the masses, conformal weights, and cubic coupling of the three asymptotic scalars. An integral formula for C involving Witten-like diagrams on the hyperbolic slices is derived. In general this integral may not be computable in closed form, but it simplifies in the near-extremal case when the incoming particle is only slightly heavier than the sum of the outgoing particles and is explicitly given below.

Other tractable examples would be of great interest. In particular, the beautiful structure of $\mathcal{N}=4$ amplitudes suggest they may take a particularly nice 2D form rewritten as correlators on \mathcal{CS}^2 . In [8] one contribution to such amplitudes (from the interior of the forward lightcone) was expressed as a Witten diagram, but to obtain the full amplitude additional harder-to-compute contributions (from outside the lightcone) must be added in. This remains an outstanding open problem.

³This generalizes the well known constraint that any QFT that can be coupled to gravity must have a local conserved stress tensor.

The utility of hyperbolic slicing was already noticed in a context with some similarity to the present one by Ashtekar and Romano in [18]. de Boer and Solodukhin initiated a program to understand flat space holography in terms of AdS holography via hyperbolic slicing in [2]. Soft theorems and aspects of scattering were hyberbolically studied in [8, 19–21]. In the context of the recent revival of the conformal bootstrap program, the linear realization of the conformal symmetry in the embedding Minkowski space has been used to simplify computations of, for example, conformal blocks and propagators in AdS [22–26].

The outline of the paper is as follows. In Section 2 we define and construct the massive and massless scalar wavefunctions that are conformal primaries of the Lorentz group $SL(2,\mathbb{C})$. Our construction, equation (2.10) below, is a convolution of plane waves with the bulk-to-boundary propagator on the hyperbolic slice H_3 , and is evaluated in terms of Bessel functions. We also present an integral transform that takes a massive scalar scattering amplitude into an $SL(2,\mathbb{C})$ covariant correlation function. In Section 3 we compute the three-point amplitude of massive conformal primary wavefunctions in the near-extremal limit. The main result is equation (3.13). In Appendix A we compute the Klein-Gordon inner product of these primary wavefunctions for a fixed mass.

2 Conformal Primary Wavefunctions

In this section we construct scalar wavefunctions that are conformal primaries of the Lorentz group $SL(2,\mathbb{C})$. A scalar conformal primary wavefunction $\phi_{\Delta,m}(X^{\mu}; w, \bar{w})$ of mass m and conformal dimension Δ is defined by the following two properties:

1. It is a solution to the Klein-Gordon equation of mass m,⁴

$$\left(\frac{\partial}{\partial X^{\nu}}\frac{\partial}{\partial X_{\nu}} - m^2\right)\phi_{\Delta,m}(X^{\mu}; w, \bar{w}) = 0.$$
(2.1)

2. It transforms covariantly as a conformal (quasi-)primary operator in two dimensions under an $SL(2,\mathbb{C})$ Lorentz transformation,

$$\phi_{\Delta,m}\left(\Lambda^{\mu}_{\nu}X^{\nu}; \frac{aw+b}{cw+d}, \frac{\bar{a}\bar{w}+\bar{b}}{\bar{c}\bar{w}+\bar{d}}\right) = |cw+d|^{2\Delta}\phi_{\Delta,m}\left(X^{\mu}; w, \bar{w}\right), \qquad (2.2)$$

where $a, b, c, d \in \mathbb{C}$ with ad - bc = 1 and Λ^{μ}_{ν} is its associated $SL(2, \mathbb{C})$ group element in the four-dimensional representation.⁵

Note that, in contrast to the situation in AdS/CFT, the mass m and the conformal dimension Δ are not related.

⁴We will use the (-,+,+,+) convention for the signature in $\mathbb{R}^{1,3}$.

⁵There is no canonical way to embed the celestial sphere into the lightcone in Minkowski space. It follows

2.1 Integral Representation

Conformal primary wavefunctions for a massive scalar field can be constructed from the Fourier transform of the bulk-to-boundary propagator in three-dimensional hyperbolic space H_3 . Let (y, z, \bar{z}) be the Poincaré coordinates of the H_3 with metric,

$$ds_{H_3}^2 = \frac{dy^2 + dz d\bar{z}}{y^2} \,. \tag{2.3}$$

Here $0 < y < \infty$ and y = 0 is the boundary of the H_3 . This geometry has an $SL(2,\mathbb{C})$ isometry that acts as

$$z \to z' = \frac{(az+b)(\bar{c}\bar{z}+\bar{d}) + a\bar{c}y^2}{|cz+d|^2 + |c|^2y^2},$$

$$\bar{z} \to \bar{z}' = \frac{(\bar{a}\bar{z}+\bar{b})(cz+d) + \bar{a}cy^2}{|cz+d|^2 + |c|^2y^2},$$

$$y \to y' = \frac{y}{|cz+d|^2 + |c|^2y^2},$$
(2.4)

with $a, b, c, d \in \mathbb{C}$ and ad - bc = 1. The H_3 can be embedded into $\mathbb{R}^{1,3}$ as either one of the two branches $(\hat{p}^0 > 0 \text{ or } \hat{p}^0 < 0)$ of the unit hyperboloid

$$-(\hat{p}^0)^2 + (\hat{p}^1)^2 + (\hat{p}^2)^2 + (\hat{p}^3)^2 = -1.$$
(2.5)

More explicitly, we can choose this embedding map $\hat{p}^{\mu}: H_3 \to \mathbb{R}^{1,3}$ for the upper hyperboloid, corresponding to an outgoing particle, to be

$$\hat{p}^{\mu}(y,z,\bar{z}) = \left(\frac{1+y^2+|z|^2}{2y}, \frac{\text{Re}(z)}{y}, \frac{\text{Im}(z)}{y}, \frac{1-y^2-|z|^2}{2y}\right). \tag{2.6}$$

This implies the useful relation

$$z = \frac{\hat{p}^1 + i\hat{p}^2}{\hat{p}^0 + \hat{p}^3} \,. \tag{2.7}$$

Let $G_{\Delta}(y, z, \bar{z}; w, \bar{w})$ be the scalar bulk-to-boundary propagator of conformal dimension Δ in H_3 [27],

$$G_{\Delta}(y, z, \bar{z}; w, \bar{w}) = \left(\frac{y}{y^2 + |z - w|^2}\right)^{\Delta}. \tag{2.8}$$

that there is also no canonical way to associate a Möbius action on w with an $SL(2,\mathbb{C})$ element Λ^{μ}_{ν} in the four-dimensional representation. In fact, any two Λ^{μ}_{ν} 's that differ by an $SL(2,\mathbb{C})$ conjugation are equally good for our definition. Below we will make a choice of the map $\Lambda^{\mu}_{\nu}(a,b,c,d)$ by fixing a reference frame for \hat{p}^{μ} in (2.6). More explicitly, Λ^{μ}_{ν} is the $SL(2,\mathbb{C})$ transformation matrix acting on \hat{p}^{μ} given by plugging (2.4) into (2.6).

This transforms covariantly under the $SL(2,\mathbb{C})$ transformation (2.4),

$$G_{\Delta}(y', z', \bar{z}'; w', \bar{w}') = |cw + d|^{2\Delta} G_{\Delta}(y, z, \bar{z}; w, \bar{w}),$$
 (2.9)

where w' = (aw + b)/(cw + d) and $\bar{w}' = (\bar{a}\bar{w} + \bar{b})/(\bar{c}\bar{w} + \bar{d})$.

The conformal primary wavefunction for a massive scalar is

$$\phi_{\Delta,m}^{\pm}(X^{\mu}; w, \bar{w}) = \int_0^\infty \frac{dy}{y^3} \int dz d\bar{z} G_{\Delta}(y, z, \bar{z}; w, \bar{w}) \exp\left[\pm im \,\hat{p}^{\mu}(y, z, \bar{z}) \,X_{\mu}\right], \qquad (2.10)$$

where we pick the minus (plus) sign for an incoming (outgoing) particle. In the next subsection we will see that potentially divergent integrals can be regulated in an $SL(2,\mathbb{C})$ covariant manner by complexifying the mass m and (2.10) is expressed in terms of Bessel functions.

It is trivial to check that (2.10) is indeed a conformal primary wavefunction. First, it satisfies the massive Klein-Gordon equation because each plane wave factor $e^{im\hat{p}\cdot X}$ does. Second, it is a conformal quasi-primary (in the sense of (2.2)) because of the $SL(2,\mathbb{C})$ covariance (2.9) of the bulk-to-boundary propagator in H_3 . Our definition and formula for the conformal primary wavefunction (2.10) can be readily generalized to Minkowski space of any dimension $\mathbb{R}^{1,d+1}$ and it would transform covariantly under the Euclidean d-dimensional conformal group SO(1, d+1).

The H_3 is embedded, via the map (2.6), into the hyperboloid in the momentum space, rather than position space and the boundary point w, \bar{w} might seem to live at the boundary of momentum space rather than Minkowski space. However, these spaces are canonically identified. The trajectory of a free massive particle is

$$X^{\mu}(s) = \hat{p}^{\mu}s + X_0^{\mu}. \tag{2.11}$$

At late times, $s \to \infty$, $-X^2 \to \infty$ and

$$\frac{X^{\mu}}{\sqrt{-X^2}} \to \hat{p}^{\mu} \,. \tag{2.12}$$

That is, massive particles asymptote to a fixed position on the hyperbolic slices of Minkowski determined by their four-momenta. Hence (w, \bar{w}) can be interpreted as a boundary coordinate of the late-time asymptotic H_3 slice.

Although we are far from constructing any example of such, the authors of [2] speculate on a boundary 2D CFT (of some exotic variety) on \mathcal{CS}^2 a subset of whose correlation functions are the 4D bulk Minkowski scattering amplitudes. Every bulk field would be dual to a continuum of operators labelled by their conformal weights. In this putative 2D CFT, the scalar bulk field mode (2.10) would be holographically dual to a local scalar boundary operator $\mathcal{O}_{\Delta}(w, \bar{w})$ of dimension Δ .

The $SL(2,\mathbb{C})$ covariance of the conformal primary wavefunction implies the $SL(2,\mathbb{C})$ covariance of any scattering amplitudes constructed from them. Let p_j^{μ} be the on-shell momenta of n massive scalars of masses m_j ($j=1,\cdots,n$). Given any Lorentz invariant n-point momentum-space scattering amplitude $\mathcal{A}(p_j^{\mu})$ of these massive scalars (including the momentum conservation delta function $\delta^{(4)}(\sum_{j=1}^n p_j^{\mu})$), the conformal primary amplitudes $\tilde{\mathcal{A}}_{\Delta_1,\cdots,\Delta_n}(w_i,\bar{w}_i)$ are

$$\widetilde{\mathcal{A}}_{\Delta_1,\dots,\Delta_n}(w_i,\bar{w}_i) \equiv \prod_{i=1}^n \int_0^\infty \frac{dy_i}{y_i^3} \int dz_i d\bar{z}_i \, G_{\Delta_i}(y_i,z_i,\bar{z}_i;w_i,\bar{w}_i) \, \mathcal{A}(m_j \hat{p}_j^{\mu}) \,,$$
 (2.13)

where $\hat{p}_j^{\mu} \equiv \hat{p}^{\mu}(y_j, z_j, \bar{z}_j)$ is given by (2.6) satisfying $\hat{p}_i^2 = -1$. By construction $\tilde{\mathcal{A}}_{\Delta_1, \dots, \Delta_n}(w_i, \bar{w}_i)$ transforms covariantly under $SL(2, \mathbb{C})$,

$$\tilde{\mathcal{A}}_{\Delta_1,\dots,\Delta_n}\left(\frac{aw_i+b}{cw_i+d},\frac{\bar{a}\bar{w}_i+\bar{b}}{\bar{c}\bar{w}_i+\bar{d}}\right) = \left(\prod_{i=1}^n |cw_i+d|^{2\Delta_i}\right)\tilde{\mathcal{A}}_{\Delta_1,\dots,\Delta_n}(w_i,\bar{w}_i). \tag{2.14}$$

2.2 Analytic Continuation and the Massless Wavefunction

The integral expression (2.10) is only a formal definition for the conformal primary wavefunction since the integral is divergent for real mass m. More rigorously, we should define our conformal primary wavefunction by analytic continuation of the integral expression (2.10) from an unphysical region. When the mass is purely imaginary $m \in -i\mathbb{R}_+$ and X^{μ} lies inside the future lightcone, the outgoing wavefunction (2.10) is convergent and can be evaluated as

$$\phi_{\Delta,m}^{+}(X^{\mu}; w, \bar{w}) = \frac{4\pi}{|m|} \frac{(\sqrt{-X^{2}})^{\Delta - 1}}{(-X^{\mu}q_{\mu})^{\Delta}} K_{\Delta - 1} \left(|m|\sqrt{-X^{2}} \right) ,$$
if $X^{0} > 0$, $X^{\mu}X_{\mu} < 0$, $m \in -i\mathbb{R}_{+}$, (2.15)

where q^{μ} is a null vector in $\mathbb{R}^{1,3}$ defined as

$$q^{\mu} = \left(1 + |w|^2, w + \bar{w}, -i(w - \bar{w}), 1 - |w|^2\right). \tag{2.16}$$

After landing on the Bessel function expression (2.15), we can then analytically continue it to real mass m and other regions in $\mathbb{R}^{1,3}$,

$$\phi_{\Delta,m}^{\pm}(X^{\mu}; w, \bar{w}) = \frac{4\pi}{im} \frac{(\sqrt{-X^2})^{\Delta - 1}}{(-X^{\mu}q_{\mu} \mp i\epsilon)^{\Delta}} K_{\Delta - 1} \left(im\sqrt{-X^2}\right). \tag{2.17}$$

We have introduced an $i\epsilon$ prescription to regularize the integral (2.10) in the case of real mass m. In practice, the integral representation (2.10) will however prove to be more convenient to compute the scattering amplitudes of these conformal primary wavefunctions.

We note that at late times inside the future lightcone, the wave equation takes the asymptotic form

$$(\partial^{\mu}\partial_{\mu} - m^2)\phi = (-\partial_{\tau}^2 - \frac{3}{\tau}\partial_{\tau} - m^2 + \cdots)\phi, \qquad \tau^2 = -X^{\mu}X_{\mu} \to \infty.$$
 (2.18)

This is solved to leading nontrivial order at large τ by

$$\phi = \frac{e^{\pm im\tau}}{\tau^{3/2}}\phi^{(0)}(y, z, \bar{z}) + \cdots, \qquad (2.19)$$

where $\phi^{(0)}$ is any function on H_3 . One may check that (2.17) with $X^2 < 0$ takes this form. On the other hand, outside the lightcone near spatial infinity we have

$$(\partial^{\mu}\partial_{\mu} - m^2)\phi = (\partial_{\sigma}^2 + \frac{3}{\sigma}\partial_{\sigma} - m^2 + \cdots)\phi, \qquad \sigma^2 = X^{\mu}X_{\mu} \to \infty.$$
 (2.20)

This is solved to leading nontrivial order at large σ by

$$\phi = \frac{e^{\pm m\sigma}}{\sigma^{3/2}}\tilde{\phi}^{(0)}(y, z, \bar{z}) + \cdots, \qquad (2.21)$$

with $\tilde{\phi}^{(0)}$ any function on dS₃. One may verify that (2.17) with $X^2 > 0$ takes this form.⁶

From the Bessel function expression (2.17) we can take the $m \to 0$ limit to obtain the massless conformal primary wavefunction (assuming $\text{Re}(\Delta) > 1$),⁷

$$\phi_{\Delta,m=0}^{\pm}(X^{\mu}; w, \bar{w}) = \frac{1}{(-X^{\mu}q_{\mu} \mp i\epsilon)^{\Delta}}.$$
 (2.22)

The massless conformal primary wavefunction has been considered in [2, 8, 19–21].

3 Massive Three-Point Amplitude

In this section we will consider the tree-level three-point amplitude $\tilde{\mathcal{A}}(w_i, \bar{w}_i)$ of the conformal primary wavefunction (2.10) $\phi_{\Delta_i, m_i}^{\pm}(X^{\mu}; w_i, \bar{w}_i)$, interacting through a local cubic vertex in $\mathbb{R}^{1,3}$,

$$\mathcal{L} \sim \lambda \phi_1 \phi_2 \phi_3 + \cdots \tag{3.1}$$

The three point scattering amplitude for plane waves is then simply

$$\mathcal{A}(p_i) = i(2\pi)^4 \lambda \,\delta^{(4)}(-p_1 + p_2 + p_3). \tag{3.2}$$

⁶One should take the square root in (2.17) corresponding to the decaying exponent when X^{μ} is outside the lightcone.

⁷Here we have dropped an overall constant factor compared to the massless limit of (2.17).

For conformal primary wavefunctions we have

$$\tilde{\mathcal{A}}(w_i, \bar{w}_i) = i\lambda \int d^4 X \,\phi_{\Delta_1, m_1}^-(X^\mu; w_1, \bar{w}_1) \prod_{i=2}^3 \phi_{\Delta_i, m_i}^+(X^\mu; w_i, \bar{w}_i) \,, \tag{3.3}$$

where we take the first particle to be incoming and the other two be outgoing. The threepoint amplitude is fixed by the $SL(2,\mathbb{C})$ covariance (2.2) to be proportional to the standard three-point function in a two-dimensional CFT:

$$\tilde{\mathcal{A}}(w_i, \bar{w}_i) \sim \frac{\lambda}{|w_1 - w_2|^{\Delta_1 + \Delta_2 - \Delta_3} |w_2 - w_3|^{\Delta_2 + \Delta_3 - \Delta_1} |w_3 - w_1|^{\Delta_3 + \Delta_1 - \Delta_2}}, \tag{3.4}$$

but it is nevertheless satisfying to see this formula explicitly appear in a 4D scattering amplitude. We wish to determine the finite proportionality constant which is a function of the masses m_i and the conformal dimensions Δ_i . In general these are given by the integral expression (3.3) which may not be possible to analytically evaluate. We will compute this constant explicitly in the near-extremal case when the mass of the first particle is slightly heavier than the sum of those of the other two. In this case the integral simplifies considerably and the three-point amplitude reduces to the tree-level three-point Witten diagram in H_3 .

Let the mass of the first particle be $2(1+\varepsilon)m$ with $\varepsilon \geq 0$ and the masses of the other two particles be m. Evaluating the X^{μ} -integral, we arrive at the following expression for the scalar three-point amplitude,⁸

$$\tilde{\mathcal{A}}(w_i, \bar{w}_i) = i(2\pi)^4 \lambda m^{-4} \left(\prod_{i=1}^3 \int_0^\infty \frac{dy_i}{y_i^3} \int dz_i d\bar{z}_i \right) \prod_{i=1}^3 G_{\Delta_i}(y_i, z_i, \bar{z}_i; w_i, \bar{w}_i) \, \delta^{(4)}(-2(1+\varepsilon)\hat{p}_1 + \hat{p}_2 + \hat{p}_3) \,,$$
(3.5)

where $\hat{p}_i^{\mu} \equiv \hat{p}^{\mu}(y_i, z_i, \bar{z}_i)$ as defined in (2.6). Note the integral does *not* take the form of a tree-level three-point Witten diagram in H_3 for general $\varepsilon \geq 0$.

We now perform the y_3, z_3, \bar{z}_3 -integrals to get rid of three delta functions. As a result, we have

$$\tilde{\mathcal{A}}(w_i, \bar{w}_i) = i2(2\pi)^4 \lambda m^{-4} \left(\int_0^\infty \frac{dy_1}{y_1^3} \int dz_1 d\bar{z}_1 \right) \left(\int_0^\infty \frac{dy_2}{y_2^3} \int dz_2 d\bar{z}_2 \right) \prod_{i=1}^3 G_{\Delta_i}(y_i, z_i, \bar{z}_i; w_i, \bar{w}_i)$$

$$\times \frac{1}{2(1+\varepsilon)y_1^{-1} - y_2^{-1}} \delta \left(\frac{2(1+\varepsilon)}{y_1 - 2(1+\varepsilon)y_2} \left[-2\varepsilon y_1 y_2 + (y_2 - y_1)^2 + |z_2 - z_1|^2 \right] \right), \tag{3.6}$$

⁸We will use the integral representation (2.10) of the conformal primary wavefunction to simplify the calculation of the amplitude and eventually make contact with the Witten diagram in H_3 in the near extremal limit. However, as discussed at the end of Section 2, the integral representation of our the conformal primary wavefunction is divergent for real mass m and the proper definition requires an analytic continuation from the Bessel function expression (2.15). Nonetheless, we will see that the three-point amplitude computed using the integral representation turns out to be finite.

where we have replaced

$$y_3 = \frac{1}{2(1+\varepsilon)y_1^{-1} - y_2^{-1}}, \qquad z_3 = \frac{2(1+\varepsilon)y_1^{-1}z_1 - y_2^{-1}z_2}{2(1+\varepsilon)y_1^{-1} - y_2^{-1}}.$$
 (3.7)

The overall factor 2 is due to the Jacobian coming from rearranging the arguments of the delta functions. Now let us perform a change of variables from (y_2, z_2, \bar{z}_2) to (R, θ, ϕ) ,

$$y_2 = y_1 + R\cos\theta$$
, $z_2 = z_1 + R\sin\theta e^{i\phi}$, $0 \le \theta \le \theta_*(y_1, R)$. (3.8)

The upper bound of θ is given by

$$\theta_*(y_1, R) = \begin{cases} \pi, & \text{if } R < y_1, \\ \cos^{-1}\left(-\frac{y_1}{R}\right), & \text{if } R \ge y_1. \end{cases}$$
 (3.9)

which comes from the constraint $\cos \theta = \frac{y_2 - y_1}{R} \ge -\frac{y_1}{R}$. We can then rewrite the three-point amplitude as

$$\tilde{\mathcal{A}}(w_i, \bar{w}_i) = i2(2\pi)^4 \lambda m^{-4} \int_0^\infty \frac{dy_1}{y_1^3} \int dz_1 d\bar{z}_1 \int_0^\infty dR R^2 \int_0^{\theta_*(y_1, R)} d\theta \sin\theta \int_0^{2\pi} d\phi \\
\times \prod_{i=1}^3 G_{\Delta_i}(y_i, z_i, \bar{z}_i; w_i, \bar{w}_i) \frac{y_1}{((2\varepsilon + 1)y_1 + 2(1+\varepsilon)R\cos\theta)(y_1 + R\cos\theta)^2} \\
\times \delta \left(\frac{2(1+\varepsilon)}{(2\varepsilon + 1)y_1 + 2(1+\varepsilon)R\cos\theta} \left[R^2 - 2\varepsilon y_1(y_1 + R\cos\theta) \right] \right), \tag{3.10}$$

with y_2, z_2, \bar{z}_2 and y_3, z_3, \bar{z}_3 replaced by (3.8) and (3.7). The delta function has support on

$$R = \sqrt{\varepsilon} \sqrt{2 + \varepsilon \cos^2 \theta} y_1 + \varepsilon y_1 \cos \theta. \tag{3.11}$$

So far we have not taken any limit on the masses $2(1+\varepsilon)m, m, m$ of the three particles. Now let us consider the near extremal limit $\varepsilon \to 0$. In this limit the three momenta $m_i\hat{p}_i$ become collinear and the corresponding points (y_i, z_i, \bar{z}_i) in H_3 become coincident. To leading order in $\sqrt{\varepsilon}$, the solution of R can be approximated by

$$R = \sqrt{2\varepsilon}y_1 + \mathcal{O}(\varepsilon). \tag{3.12}$$

In the near extremal limit we have $R \sim 0$, hence the three bulk points y_i, z_i, \bar{z}_i in H_3 become coincident as commented above. We can then bring the three bulk-to-boundary propagators outside the R, θ, ϕ -integral. Next, by a simple power counting of R, we find that the three-point amplitude of the conformal primary wavefunction is zero when $\varepsilon = 0$, which is related to the fact that the phase space vanishes for marginal decay process. We should proceed

to the subleading order in the near extremal expansion to obtain a nonzero answer, which is⁹

$$\tilde{\mathcal{A}}(w_{i}, \bar{w}_{i}) = \frac{i(2\pi)^{5}\lambda}{m^{4}} \int_{0}^{\infty} \frac{dy_{1}}{y_{1}^{3}} \int dz_{1} d\bar{z}_{1} \prod_{i=1}^{3} G_{\Delta_{i}}(y_{1}, z_{1}, \bar{z}_{1}; w_{i}, \bar{w}_{i})
\times \frac{1}{y_{1}} \int_{0}^{\infty} dR R^{2} \int_{0}^{\pi} d\theta \sin \theta \, \delta \left(R^{2} - 2\varepsilon y_{1}^{2}\right) + \mathcal{O}(\varepsilon)
= \frac{i2^{\frac{11}{2}}\pi^{5}\lambda}{m^{4}} \sqrt{\varepsilon} \left(\int_{0}^{\infty} \frac{dy_{1}}{y_{1}^{3}} \int dz_{1} d\bar{z}_{1} \prod_{i=1}^{3} G_{\Delta_{i}}(y_{1}, z_{1}, \bar{z}_{1}; w_{i}, \bar{w}_{i}) \right) + \mathcal{O}(\varepsilon)
= \frac{i2^{\frac{9}{2}}\pi^{6}\lambda \Gamma(\frac{\Delta_{1} + \Delta_{2} + \Delta_{3} - 2}{2}) \Gamma(\frac{\Delta_{1} + \Delta_{2} - \Delta_{3}}{2}) \Gamma(\frac{\Delta_{1} - \Delta_{2} + \Delta_{3}}{2}) \Gamma(\frac{-\Delta_{1} + \Delta_{2} + \Delta_{3}}{2}) \sqrt{\varepsilon}}{m^{4}\Gamma(\Delta_{1})\Gamma(\Delta_{2})\Gamma(\Delta_{3}) |w_{1} - w_{2}|^{\Delta_{1} + \Delta_{2} - \Delta_{3}} |w_{2} - w_{3}|^{\Delta_{2} + \Delta_{3} - \Delta_{1}} |w_{3} - w_{1}|^{\Delta_{3} + \Delta_{1} - \Delta_{2}}} + \mathcal{O}(\varepsilon),$$

where the term in the parentheses in the second to last line is precisely the tree-level three-point Witten diagram in H_3 , which was evaluated in [28]. Hence the near extremal massive three-point amplitude takes the form of the three-point function of scalar primaries with conformal dimensions Δ_i in a two-dimensional CFT.

Acknowledgements

We are grateful to T. Dumitrescu, P. Mitra, M. Pate, B. Schwab, D. Simmons-Duffin, and A. Zhiboedov for useful conversations. This work was supported in part by NSF grant 1205550. S.P. is supported by the NSF and by the Hertz Foundation through a Harold and Ruth Newman Fellowship. S.H.S. is supported by the National Science Foundation grant PHY-1606531.

A Klein-Gordon Inner Product

In this section we evaluate the Klein-Gordon inner product between two conformal primary solutions with the same mass m and generic complex weights $\Delta_{1,2}$. $SL(2,\mathbb{C})$ implies this must vanish for $\Delta_1 \neq \Delta_2^*$, while some kind of delta function is expected at $\Delta_1 = \Delta_2^*$.

The Klein-Gordon inner product between two outgoing wavefunctions $\phi_{\Delta_1,m}^+(X^\mu;w_1,\bar{w}_1)$

⁹Note that in the near extremal limit $\varepsilon \to 0$, the upper bound $\theta_*(y_1, R)$ of the angular coordinate becomes π on the support of the delta function.

and $\phi_{\Delta_2,m}^+(X^\mu; w_2, \bar{w}_2)$ evaluated on the slice $X^0=0$ is

$$(\phi_{1}^{+},\phi_{2}^{+}) = -i \int d^{3}\vec{X} \left[\phi_{\Delta_{1},m}^{+}(X^{\mu}; w_{1}, \bar{w}_{1}) \partial_{X^{0}} \phi_{\Delta_{2},m}^{+*}(X^{\mu}; w_{2}, \bar{w}_{2}) \right. \\ \left. - \partial_{X^{0}} \phi_{\Delta_{1},m}^{+}(X^{\mu}; w_{1}, \bar{w}_{1}) \phi_{\Delta_{2},m}^{+*}(X^{\mu}; w_{2}, \bar{w}_{2}) \right] \\ = (2\pi)^{3} m^{-2} \left(\prod_{i=1}^{2} \int_{0}^{\infty} \frac{dy_{i}}{y_{i}^{3}} \int dz_{i} d\bar{z}_{i} \right) G_{\Delta_{1}}(y_{1}, z_{1}, \bar{z}_{1}; w_{1}, \bar{w}_{1}) G_{\Delta_{2}}^{*}(y_{2}, z_{2}, \bar{z}_{2}; w_{2}, \bar{w}_{2}) \\ \times \left(\frac{1 + y_{1}^{2} + |z_{1}|^{2}}{2y_{1}} + \frac{1 + y_{2}^{2} + |z_{2}|^{2}}{2y_{2}} \right) \delta^{(2)} \left(\frac{z_{1}}{y_{1}} - \frac{z_{2}}{y_{2}} \right) \delta \left(\frac{1 - y_{1}^{2} - |z_{1}|^{2}}{2y_{1}} - \frac{1 - y_{2}^{2} - |z_{2}|^{2}}{2y_{2}} \right) \\ = 2(2\pi)^{3} m^{-2} \int_{0}^{\infty} \frac{dy}{y^{3}} \int dz d\bar{z} G_{\Delta_{1}}(y, z, \bar{z}; w_{1}, \bar{w}_{1}) G_{\Delta_{2}}^{*}(y, z, \bar{z}; w_{2}, \bar{w}_{2}) . \tag{A.1}$$

Using the Feynman trick,

$$\frac{1}{A^a B^b} = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \int_0^1 d\alpha \frac{\alpha^{a-1} (1-\alpha)^{b-1}}{(\alpha A + (1-\alpha)B)^{a+b}}, \tag{A.2}$$

we can perform the integrals in y, z, \bar{z} to obtain

$$(\phi_1^+, \phi_2^+) = 2(2\pi)^3 m^{-2} \frac{\pi \Gamma(\frac{\Delta_1 + \Delta_2^* - 2}{2}) \Gamma(\frac{\Delta_1 + \Delta_2^*}{2})}{2\Gamma(\Delta_1) \Gamma(\Delta_2^*) |w_1 - w_2|^{\Delta_1 + \Delta_2^*}} \int_0^1 d\alpha \alpha^{\frac{\Delta_1 - \Delta_2^*}{2} - 1} (1 - \alpha)^{\frac{-\Delta_1 + \Delta_2^*}{2} - 1}.$$
 (A.3)

Here Δ_2^* is the complex conjugate of Δ_2 . If we let $\eta = \frac{\Delta_1 - \Delta_2^*}{2}$, $\alpha = \frac{e^u}{e^u + e^{-u}}$

$$\int_0^1 d\alpha \alpha^{\eta - 1} (1 - \alpha)^{-\eta - 1} = 2 \int_{-\infty}^\infty du e^{2u\eta} . \tag{A.4}$$

This is divergent if η is real, and equals to $2\pi\delta(\lambda)$ if $\eta=i\lambda$ is pure imaginary. Therefore, in order to have a delta function normalizable inner product, we require Δ_i 's to be complex numbers with the same real part, $\Delta_1 = a + i\lambda_1$, $\Delta_2 = a + i\lambda_2$ $(a, \lambda_i \in \mathbb{R})$. The Klein-Gordon inner product for complex conformal dimensions, equal mass conformal primary wavefunction is,

$$(\phi_1^+, \phi_2^+) = 64\pi^5 m^{-2} \frac{1}{(\Delta_1 + \Delta_2^* - 2) |w_1 - w_2|^{\Delta_1 + \Delta_2^*}} \delta(\lambda_1 + \lambda_2). \tag{A.5}$$

References

[1] P. A. M. Dirac, "Wave equations in conformal space," Annals Math. 37 (1936) 429–442.

- [2] J. de Boer and S. N. Solodukhin, "A Holographic reduction of Minkowski space-time," Nucl. Phys. **B665** (2003) 545–593, hep-th/0303006.
- [3] T. Banks, "A Critique of pure string theory: Heterodox opinions of diverse dimensions," hep-th/0306074.
- [4] G. Barnich and C. Troessaert, "Symmetries of asymptotically flat 4 dimensional spacetimes at null infinity revisited," *Phys. Rev. Lett.* **105** (2010) 111103, 0909.2617.
- [5] G. Barnich and C. Troessaert, "Supertranslations call for superrotations," *PoS* (2010) 010, 1102.4632. [Ann. U. Craiova Phys.21,S11(2011)].
- [6] D. Kapec, V. Lysov, S. Pasterski, and A. Strominger, "Semiclassical Virasoro symmetry of the quantum gravity S-matrix," JHEP 08 (2014) 058, 1406.3312.
- [7] D. Kapec, P. Mitra, A.-M. Raclariu, and A. Strominger, "A 2D Stress Tensor for 4D Gravity," 1609.00282.
- [8] C. Cheung, A. de la Fuente, and R. Sundrum, "4D Scattering Amplitudes and Asymptotic Symmetries from 2D CFT," 1609.00732.
- [9] S. He, Y.-t. Huang, and C. Wen, "Loop Corrections to Soft Theorems in Gauge Theories and Gravity," *JHEP* 12 (2014) 115, 1405.1410.
- [10] M. Bianchi, S. He, Y.-t. Huang, and C. Wen, "More on Soft Theorems: Trees, Loops and Strings," Phys. Rev. D92 (2015), no. 6, 065022, 1406.5155.
- [11] Z. Bern, S. Davies, and J. Nohle, "On Loop Corrections to Subleading Soft Behavior of Gluons and Gravitons," *Phys. Rev.* **D90** (2014), no. 8, 085015, 1405.1015.
- [12] Z. Bern, S. Davies, P. Di Vecchia, and J. Nohle, "Low-Energy Behavior of Gluons and Gravitons from Gauge Invariance," *Phys. Rev.* **D90** (2014), no. 8, 084035, 1406.6987.
- [13] S. W. Hawking, M. J. Perry, and A. Strominger, "Superrotation Charge and Supertranslation Hair on Black Holes," 1611.09175.
- [14] T. He, D. Kapec, A.-M. Raclariu, and A. Strominger, "Loop-Corrected Virasoro Symmetry of 4D Quantum Gravity," to appear.
- [15] F. Cachazo and A. Strominger, "Evidence for a New Soft Graviton Theorem," 1404.4091.
- [16] A. Strominger, "Asymptotic Symmetries of Yang-Mills Theory," JHEP 07 (2014) 151, 1308.0589.

- [17] T. He, P. Mitra, and A. Strominger, "2D Kac-Moody Symmetry of 4D Yang-Mills Theory," *JHEP* **10** (2016) 137, 1503.02663.
- [18] A. Ashtekar and J. D. Romano, "Spatial infinity as a boundary of space-time," *Class. Quant. Grav.* **9** (1992) 1069–1100.
- [19] M. Campiglia and A. Laddha, "Asymptotic symmetries of QED and Weinberg's soft photon theorem," *JHEP* **07** (2015) 115, 1505.05346.
- [20] M. Campiglia and A. Laddha, "Asymptotic symmetries of gravity and soft theorems for massive particles," *JHEP* 12 (2015) 094, 1509.01406.
- [21] M. Campiglia, "Null to time-like infinity Green's functions for asymptotic symmetries in Minkowski spacetime," *JHEP* **11** (2015) 160, 1509.01408.
- [22] L. Cornalba, M. S. Costa, and J. Penedones, "Deep Inelastic Scattering in Conformal QCD," *JHEP* **03** (2010) 133, 0911.0043.
- [23] S. Weinberg, "Six-dimensional Methods for Four-dimensional Conformal Field Theories," Phys. Rev. D82 (2010) 045031, 1006.3480.
- [24] M. S. Costa, J. Penedones, D. Poland, and S. Rychkov, "Spinning Conformal Correlators," *JHEP* 11 (2011) 071, 1107.3554.
- [25] M. S. Costa, J. Penedones, D. Poland, and S. Rychkov, "Spinning Conformal Blocks," JHEP 11 (2011) 154, 1109.6321.
- [26] D. Simmons-Duffin, "Projectors, Shadows, and Conformal Blocks," JHEP 04 (2014) 146, 1204.3894.
- [27] E. Witten, "Anti-de Sitter space and holography," Adv. Theor. Math. Phys. 2 (1998) 253–291, hep-th/9802150.
- [28] D. Z. Freedman, S. D. Mathur, A. Matusis, and L. Rastelli, "Correlation functions in the CFT(d) / AdS(d+1) correspondence," Nucl. Phys. B546 (1999) 96–118, hep-th/9804058.